

SIMPSON TYPE INEQUALITIES FOR Q - CLASS FUNCTIONS

M.EMIN ÖZDEMİR[♦], ALPER EKINCI[♣], MUSTAFA GÜRBÜZ[♣],
AND AHMET OCAK AKDEMİR[♣]

ABSTRACT. In this paper, we obtain some Simpson type inequalities for functions whose second derivatives' absolute value or q -th power of them are Q -class functions. Also we give applications to numerical integration.

1. INTRODUCTION

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a four times continuously differentiable mapping on (a, b) and $\|f^{(4)}\|_{\infty} = \sup |f^{(4)}(x)| < \infty$. The following inequality

$$\left| \frac{1}{3} \left[\frac{f(a) + f(b)}{2} + 2f\left(\frac{a+b}{2}\right) \right] - \frac{1}{b-a} \int_a^b f(x)dx \right| \leq \frac{1}{2880} \|f^{(4)}\|_{\infty} (b-a)^4$$

is well known in the literature as Simpson's inequality.

For some results about Simpson inequality see [1]-[6].

Recall the Definition of Godunova-Levin function as following:

Definition 1. (See [7], [8, p.410]) We say that $f : I \rightarrow \mathbb{R}$ is a Godunova-Levin function or that f belongs to the class $Q(I)$ if f is non-negative and for all $x, y \in I$ and $t \in (0, 1)$ we have

$$f(tx + (1-t)y) \leq \frac{f(x)}{t} + \frac{f(y)}{1-t}.$$

The main aim of this paper is to give some new inequalities of Simpson's type for functions whose second derivatives of absolute values are Godunova-Levin functions or belong to the class $Q(I)$.

2. MAIN RESULTS

We used the following Lemma to obtain our main results.

Lemma 1. (See [6]) Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be an absolutely continuous mapping on I^0 where $a, b \in I$ with $a < b$, such that $f'' \in L[a, b]$. Then the following equality holds:

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(x)dx - \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \right| \\ &= (b-a)^2 \int_0^1 p(t) f''(tb + (1-t)a) dt, \end{aligned}$$

Key words and phrases. Simpson inequality, Q -Class functions, power-mean inequality, Simpson's formulae.

where

$$p(t) = \begin{cases} \frac{1}{6}t(3t-1), & t \in [0, \frac{1}{2}) \\ \frac{1}{6}(t-1)(3t-2), & t \in [\frac{1}{2}, 1] \end{cases}.$$

Theorem 1. *Let f be a functions which satisfy the assumptions of Lemma 1. If $|f''|$ belongs to the class $Q(I)$, then the following inequality holds:*

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(x)dx - \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \right| \\ & \leq \frac{(b-a)^2}{6} \left(\frac{12 \ln 2 - 8 \ln 3 + 1}{2} \right) [|f''(a)| + |f''(b)|]. \end{aligned}$$

Proof. From Lemma 1 and using the properties of modulus, we have

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(x)dx - \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \right| \\ & \leq (b-a)^2 \left\{ \int_0^{\frac{1}{2}} \left| \frac{1}{6}t(3t-1) \right| |f''(tb + (1-t)a)| dt \right. \\ & \quad \left. + \int_{\frac{1}{2}}^1 \left| \frac{1}{6}(t-1)(3t-2) \right| |f''(tb + (1-t)a)| dt \right\} \\ & = \frac{(b-a)^2}{6} \left\{ \int_0^{\frac{1}{3}} t(1-3t) |f''(tb + (1-t)a)| dt + \int_{\frac{1}{3}}^{\frac{1}{2}} t(3t-1) |f''(tb + (1-t)a)| dt \right. \\ & \quad \left. + \int_{\frac{1}{2}}^{\frac{2}{3}} (1-t)(2-3t) |f''(tb + (1-t)a)| dt + \int_{\frac{2}{3}}^1 (1-t)(3t-2) |f''(tb + (1-t)a)| dt \right\}. \end{aligned}$$

Since $|f''|$ belongs to the class $Q(I)$, we can write

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(x)dx - \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \right| \\ & \leq \frac{(b-a)^2}{6} \left\{ \int_0^{\frac{1}{3}} t(1-3t) \left(\frac{|f''(b)|}{t} + \frac{|f''(a)|}{1-t} \right) dt + \int_{\frac{1}{3}}^{\frac{1}{2}} t(3t-1) \left(\frac{|f''(b)|}{t} + \frac{|f''(a)|}{1-t} \right) dt \right. \\ & \quad \left. + \int_{\frac{1}{2}}^{\frac{2}{3}} (1-t)(2-3t) \left(\frac{|f''(b)|}{t} + \frac{|f''(a)|}{1-t} \right) dt + \int_{\frac{2}{3}}^1 (1-t)(3t-2) \left(\frac{|f''(b)|}{t} + \frac{|f''(a)|}{1-t} \right) dt \right\} \end{aligned}$$

Computing the above integrals, we get the desired result. $\square$

Corollary 1. *In Theorem 1, if we take $f(a) = f(b) = f\left(\frac{a+b}{2}\right)$, then we have*

$$\left| \frac{1}{b-a} \int_a^b f(x)dx - f\left(\frac{a+b}{2}\right) \right| \leq \frac{(b-a)^2}{6} \left(\frac{12 \ln 2 - 8 \ln 3 + 1}{2} \right) [|f''(a)| + |f''(b)|].$$

For $M > 0$, if $|f''(x)| < M$, for all $x \in [a, b]$, then we have

$$\left| \frac{1}{b-a} \int_a^b f(x)dx - f\left(\frac{a+b}{2}\right) \right| \leq \frac{(b-a)^2}{3} \left(\frac{12 \ln 2 - 8 \ln 3 + 1}{2} \right) M.$$

Theorem 2. Let f be a functions which satisfy the assumptions of Lemma 1. If $|f''|^q$ belongs to the class $Q(I)$, then the following inequality holds:

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(x) dx - \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \right| \\ & \leq \frac{(b-a)^2}{6} \left(\frac{1}{27} \right)^{1-\frac{1}{q}} \left\{ \left[\left(6 \ln 2 - 4 \ln 3 + \frac{7}{24} \right) |f''(a)|^q + \frac{5|f''(b)|^q}{24} \right]^{\frac{1}{q}} \right. \\ & \quad \left. + \left[\left(6 \ln 2 - 4 \ln 3 + \frac{7}{24} \right) |f''(b)|^q + \frac{5|f''(a)|^q}{24} \right]^{\frac{1}{q}} \right\}. \end{aligned}$$

where $q \geq 1$.

Proof. From Lemma 1 and using the Power-mean inequality, we have

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(x) dx - \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \right| \\ & \leq \frac{(b-a)^2}{6} \left\{ \left(\int_0^{\frac{1}{2}} |t(3t-1)| dt \right)^{1-\frac{1}{q}} \left(\int_0^{\frac{1}{2}} |t(3t-1)| |f''(tb + (1-t)a)|^q dt \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\int_{\frac{1}{2}}^1 |(t-1)(3t-2)| dt \right)^{1-\frac{1}{q}} \left(\int_{\frac{1}{2}}^1 |(t-1)(3t-2)| |f''(tb + (1-t)a)|^q dt \right)^{\frac{1}{q}} \right\} \end{aligned}$$

Since $|f''|$ belongs to the class $Q(I)$ and by computing the above integrals, we deduce

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(x) dx - \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] \right| \\ & \leq \frac{(b-a)^2}{6} \left(\frac{1}{27} \right)^{1-\frac{1}{q}} \left\{ \left(\left(6 \ln 2 - 4 \ln 3 + \frac{7}{24} \right) |f''(a)|^q + \frac{5|f''(b)|^q}{24} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\left(6 \ln 2 - 4 \ln 3 + \frac{7}{24} \right) |f''(b)|^q + \frac{5|f''(a)|^q}{24} \right)^{\frac{1}{q}} \right\} \end{aligned}$$

which is the desired result. $\square$

3. APPLICATIONS TO NUMERICAL INTEGRATION

Let d be a division of the interval $[a, b]$, i.e., $d : a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$, $h_i = \frac{(x_{i+1} - x_i)}{2}$ and consider the Simpson's formulae

$$S(f, d) = \sum_{i=0}^{n-1} \frac{f(x_i) + f(x_i + h_i) + f(x_{i+1})}{6} (x_{i+1} - x_i).$$

It is well-known that if the mapping $f : [a, b] \rightarrow \mathbb{R}$, is differentiable such that $f^{(4)}(x)$ exists on (a, b) and $M = \max_{x \in (a, b)} |f^{(4)}(x)| < \infty$, then

$$I = \int_a^b f(x) dx = S(f, d) + E_s(f, d),$$

where the approximation error $E_s(f, d)$ of the integral I by the Simpson's formulae $S(f, d)$ satisfies

$$E_s(f, d) \leq \frac{M}{90} \sum_{i=0}^{n-1} (x_{i+1} - x_i)^5.$$

Now we will give estimation for remainder term $E(f, d)$ in terms of the second derivative.

Proposition 1. *Under the conditions of Theorem 1, then for every division d of $[a, b]$, the following holds:*

$$|E_s(f, d)| \leq \left(\frac{12 \ln 2 - 8 \ln 3 + 1}{12} \right) \sum_{i=0}^{n-1} (x_{i+1} - x_i)^2 [|f''(x_i)| + |f''(x_{i+1})|].$$

Proof. Applying Theorem 1 on the subintervals $[x_i, x_{i+1}]$, $(i = 0, 1, \dots, n-1)$ of the division d and summing over i from 0 to $n-1$, we get the required result. $\square$

REFERENCES

- [1] M. Alomari, M. Darus and S.S. Dragomir, New inequalities of Simpson's type for s -convex functions with applications, RGMIA Res. Rep. Coll., 12 (4) (2009).
- [2] M. Alomari and M. Darus, On some inequalities of Simpson-type via quasi-convex functions and applications, Transylvanian Journal of Mathematics and Mechanics, vol. 2, no. 1, pp. 15–24, 2010.
- [3] M.Z. Sarikaya, E. Set and M.E. Özdemir, On new inequalities of Simpson's type for s -convex functions, Comput. Math. Appl. 60 (2010) 2191–2199.
- [4] S. S. Dragomir, R. P. Agarwal, and P. Cerone, On Simpson's inequality and applications, Journal of Inequalities and Applications, vol. 5, no. 6, pp. 533–579, 2000.
- [5] N.Ujević, Sharp inequalities of Simpson type and Ostrowski type, Comp. Math. Appl., 48 (2004), 145–151.
- [6] N.Ujević, Two sharp inequalities of Simpson type and applications, Georgian Math. J., 1 (11) (2004), 187–194.
- [7] E.K. Godunova, V.I. Levin, Neravenstva dlja funkciï širokogo klassa, soderžašcego vypuklye, monotonnnye i nekotorye drugie vidy funkciï, in: Vycislitel. Mat. i. Mat. Fiz. Mežvuzov. Sb. Nauc. Trudov, MGPI, Moskva, 1985, 138–142.
- [8] D.S. Mitrinović, J. Pečarić and A.M. Fink, Classical and new inequalities in analysis, *Kluwer Academic Publishers*, Dordrecht, 1993.

[♦]ATATÜRK UNIVERSITY, K.K. EDUCATION FACULTY, DEPARTMENT OF MATHEMATICS, ERZURUM 25240, TURKEY

E-mail address: emos@atauni.edu.tr

E-mail address: alperekinci@hotmail.com

[♣]AĞRI İBRAHİM ÇEÇEN UNIVERSITY, FACULTY OF EDUCATION, DEPARTMENT OF MATHEMATICS, AĞRI 04100, TURKEY

E-mail address: mgurbuz@agri.edu.tr

[♣]AĞRI İBRAHİM ÇEÇEN UNIVERSITY, FACULTY OF SCIENCE AND ARTS, DEPARTMENT OF MATHEMATICS, AĞRI 04100, TURKEY

E-mail address: ahmetakdemir@agri.edu.tr